Magnetized Prestellar Core Formation in Turbulent Cloud

Che-Yu Chen

with Eve Ostriker
Lee Mundy
Shaye Storm
and The CLASSy Collaboration

ESA-ESTEC, Nov 11, 2014

Star Formation Theory

- A general picture



Star Formation Theory

- A general picture



Core Formation Theory

- From diffuse clouds to dense cores



Gravity
vs.
Magnetic field
+
Turbulence

self-
gravitating

$t_{core} \sim 2\text{-}5\ t_{ff}$
~ 1 Myr

Core Formation Theory

- How do magnetically supercritical cores form?

➢ Standard scenario of core evolution

quasi-static ambipolar diffusion



$t \sim 10^7$ yr

self-gravitating core

☹ core lifetime too long

➢ New model: turbulence-accelerated, magnetically-regulated core formation (Li & Nakamura 2004)

Simulation Setup



- Include:
  - self-gravity
  - perturbation field
  - ambipolar diffusion
- cloud density:
  - $n_0 = 1000\ \mathrm{cm}^{-3}$



Structure Formation



Kinematics



(Mundy et al. *in prep.*)

Line-of-sight Velocity



(Mundy et al. *in prep.*)

Filament Structure





turbulence vs. self-gravity

(Mundy et al. *in prep.*)

Observations

- CLASSy results







(Mundy et al. *in prep.*)

Filament ⇨ Core Formation

- Which one forms first?

  ⇒ filament fragmentation vs. simultaneity



- How do magnetically supercritical cores form?

Cores in Simulations



(Chen & Ostriker 2014)

Cores in Simulations



(Chen & Ostriker 2014)

Statistical Results

- Models with same pre-shock pressure
  ⇒ similar core masses, with or without AD





(Chen & Ostriker 2014)

Anisotropic Gas Flows



(Chen & Ostriker 2014)



Anisotropic Core Formation

filament formed in post-shock layer



Magnetic critical length

$$L_{\rm mag} = \frac{B_{\rm ps}}{\rho_{\rm ps}} \frac{1}{2\pi\sqrt{G}}$$

contract longitudinally until $L' \sim 2R$ and $\rho' = \frac{L}{2R}\rho_{\rm ps}$

Thermally supercritical: $R \sim R_{\rm BE}\left(\rho'\right) = 0.65\frac{c_s}{\sqrt{G\rho'}}$ ➡ $R = 0.84\frac{{c_s}^2}{G\rho_{\rm ps}L}$,

$$M = \pi R^2 L \rho_{\rm ps} = 2.2\frac{{c_s}^4}{G^2\rho_{\rm ps}L}$$

Magnetically critical, Anisotropic

$$L = L_{\rm mag} \Rightarrow \rho_{\rm ps}L = \frac{B_{\rm ps}}{2\pi\sqrt{G}}$$

$$M_{\rm crit,cyl} = 14\frac{{c_s}^4}{\sqrt{G^3}B_{\rm ps}}$$

$$= 1.3\ {\rm M}_\odot \left(\frac{B_{\rm ps}}{50\ \mu{\rm G}}\right)^{-1}\left(\frac{T}{10\ {\rm K}}\right)^2$$

$$= 2.8\frac{c_s}{\sqrt{G^3\rho_0{v_0}^2}} \propto \mathcal{M}^{-1}$$

$B_{ps}^2/8\pi \sim \rho_0 v_0^2$

Magnetically critical, Isotropic

Spherical core with $R = \frac{L_{\rm mag}}{2} = \frac{B_{\rm ps}}{4\pi\sqrt{G}\rho_{\rm ps}}$

$$M_{\rm crit,sph} = \frac{4\pi R^3}{3}\rho_{\rm ps} = \frac{1}{48\pi^2 G^{3/2}}\frac{{B_{\rm ps}}^3}{{\rho_{\rm ps}}^2}$$

$$= 5.21\ {\rm M}_\odot \left(\frac{B_{\rm ps}}{50\ \mu{\rm G}}\right)^3\left(\frac{n_{\rm ps}}{10^4\ {\rm cm}^{-3}}\right)^{-2}$$

Further Tests

- Core mass varies with inflow Mach number

$M_{\rm core} \propto \mathcal{M}^{-0.9}$

(Chen & Ostriker *in prep.*)





- $B_{\rm core}$ is within a factor of 2 of $B_{\rm post\text{-}shock}$

Summary

- Filament transverse velocity gradients provide evidence of condensation in flattened structures

- Ambipolar diffusion is not necessary for core formation, and magnetization is not the main factor controlling core mass & size

- Magnetically-supercritical, low-mass cores form anisotropically via contraction along $\vec{B}$

- These cores have masses and magnetic fields that depend on pre-shock ρv^2 in cloud